

# Prevalence Of Mastitis In Dairy Cows In Selected Areas Of Sylhet District

**Dr. Md. Matiur Rahman, Jannatul Mawa Momu, Mahfuz Rahman Adnan, Mowdudul Hasan Talha, Mamun Ur Rahman, Mustaq Ahmad, Akash Khasnobish, Md. abdu Sabur, Md Altafur Rahman**

Assistant Professor, Department of Medicine, Sylhet Agricultural University,  
Sylhet, Bangladesh.  
*matiur.dvm@sau.ac.bd*

Faculty of Veterinary, Animal and Biomedical Sciences, Sylhet Agricultural University  
MS Fellow, Department of Animal Nutrition, Bangladesh, +8801670761782  
*jannatulmumu.sau@gmail.com*

Faculty of Veterinary, Animal and Biomedical Sciences, Sylhet Agricultural University  
MS Fellow, Department of Pathology, Bangladesh, +8801746798499  
*mahfuzradnan@gmail.com*

Faculty of Veterinary, Animal and Biomedical Sciences, Sylhet Agricultural University  
MS Fellow, Department of Pharmacology and Toxicology, Bangladesh, +8801 01632860598  
*hasantalhavet@gmail.com*

Faculty of Veterinary, Animal and Biomedical Sciences, Sylhet Agricultural University  
MS Fellow, Department of Genetics and Animal Breeding, Bangladesh, +8801740202028  
*marshal7bd@gmail.com*

Faculty of Veterinary, Animal and Biomedical Sciences, Sylhet Agricultural University  
MS Fellow, Department of Genetics and Animal Breeding, Bangladesh, +8801 01632860598  
*mustaq1201005@gmail.com*

Faculty of Veterinary, Animal and Biomedical Sciences, Sylhet Agricultural University  
MS Fellow, Department of Pharmacology and Toxicology, Bangladesh, +8801746863844  
*khasnobish1201021@gmail.com*

Faculty of Veterinary, Animal and Biomedical Sciences, Sylhet Agricultural University  
MS Fellow, Department of Medicine, Bangladesh, +8801720184665  
*mdsabur06@gmail.com*

Faculty of Veterinary, Animal and Biomedical Sciences, Sylhet Agricultural University  
MS Fellow, Department of Genetics and Animal Breeding, Bangladesh, +8801712307719  
*altafrussell@gmail.com*

**ABSTRACT:** This study was carried out to determine the prevalence of mastitis in lactating Dairy Cow at District Dairy Farms in Sylhet and UVH, at Jaintapur during the period of January 2016 to December 2016. Mastitis was most frequently seen in cross breed cows. In Jaintapur Upazilla, most of cows were indigenous, but there were many farmers rear cross breed cow for the purpose of selling milk in market. In Dairy Farm, all cows were cross breed. The overall prevalence was higher in crossbred cows than indigenous cows. The prevalence of mastitis was higher in advancing age and in lactation period. The cows were more prone to mastitis during 1st month of lactation. There was significant relationship between prevalence of mastitis and associated risk factors like general physical condition, herd size, frequency of dung removal, floor drainage quality and condition of floor. Reproductive diseases and periparturiant diseases showed higher incidence of mastitis in cows. The overall prevalence of mastitis at Jaintapur was 7.15%. The prevalence of mastitis of dairy cows in Jaintapur Upazilla was more in rainy season and less in winter season. The prevalence of mastitis was 3.47% in summer season, 11.52% in rainy season and 5% in winter season. The overall prevalence of mastitis at Dairy farm was 6.91%. The prevalence of mastitis in dairy farm was more in rainy season and less in summer season. The prevalence of mastitis was 4.97% in summer season; 16.53% in rainy season and 5.29% in winter season.

**Key word:** Breed, lactating, Mastitis, Sylhet, Udder

## 1. Introduction

Bangladesh has 24 million cattle, out of which 6 million are dairy cattle of local and crossbreds [1]. The majority of the dairy cattle are in the hands of smallholder dairy producers. The estimated numbers of dairy farms in Bangladesh is 1.4 million [2]. It is the part of the mixed farming systems and a predominant source of income and nutrition and jobs. Milk is the major sources of money income from dairying. Annual milk production in

Bangladesh is 16.2 lack metric tons and about 64% milk comes from cattle [3]. However, Bangladesh has an acute shortage of milk. The produced milk can fulfill only 13.6% of the total requirement in Bangladesh. Dairying in Bangladesh is growing faster but it also faces lot of problems of high input and low output prices. Disease, along with non-availability of feed resources and nutrition are the most important constraints to milk production. The term "Mastitis" derives from Greck word Mastos which

means breast (mammary gland) and its meaning inflammation, which is inflammation of the mammary gland is called Mastitis. Mastitis is the term, which means inflammation of parenchyma of the mammary gland (udder) regardless of the cause. It is characterized by physical, chemical and microbiological changes in the milk and by pathological change in the glandular tissues of the udder. The most important changes in the milk include discoloration, the presence of clots and the presence of abnormally large number of leukocytes. Bovine mastitis (both clinical and subclinical) is one of the most important diseases causing considerable economical loss to the farmers and the dairy industry and is mainly caused by bacterial infection in the udder. The mastitis can be diagnosed on history and clinical findings but laboratory examinations ascertain the subclinical mastitis. The loss due to subclinical mastitis (SCM) has been estimated to be Rs. 1058 million per annum in India which was three times more than that caused clinical mastitis [4]. The prevalence of SCM has been shown to be 15 to 40 times more than the clinical mastitis [5]. However, no such estimation of economic loss due to mastitis is available from Bangladesh. Reduction in the milk in SCM is not only responsible for great economic loss to the dairy industry but also acts as a carrier and source of infection for healthy milch cows which is one of the biggest obstacles in the achievement of self-sufficiency in milk production in Bangladesh. Mastitis not only affects the milk yield adversely, but in many cases alters the consumption of milk in such a way that it is rendered unsuitable for preparation of certain milk products [6]. The residual effects of many commonly used antibiotics, and chemotherapeutic agents in treating cases of mastitis, pathogenic microorganism in milk (if used without boiling) can also be a potential danger to human health. So to overcome a major hurdle in the way of achieving self-sufficiency in milk production, it seems to be prime importance to prevent the incidence of mastitis by early diagnosis of subclinical form of the disease as otherwise this subclinical form may lead to clinical which is irreversible in most cases. The mastitis causative organisms are ubiquitous in nature and persist long time in the cow yard or barns and there is chance of constant udder infection under poor hygienic and management systems [7,8]. Epidemiological studies on mastitis revealed that mastitogenic organisms are widespread on different body sites of the cows, milker's hands, milking

## 2.6 Study design

$$\text{Prevalence} = \frac{\text{Number of animals with the disease}}{\text{Number of animals at risk}} \times 100$$

Percentage was used to express the prevalence of mastitis of dairy cows in Jaintapur Upazilla and Dairy Farm, Sylhet.

## 2.7 Diagnostic Procedure

Diagnosis of clinical mastitis performed on the basis of clinical signs showed by the animals. It can be interpreted in following forms-

### ➤ Mild Form

Flakes and clots were found in milk. Slight swelling of infected quarter was found. Systemic reaction and fever was absent.

cans and in the milk samples. Teat apices have been reported to be the most common site [9], from where these organisms have been isolated [10,11]. The mastitis causing organism, Staphylococci, the chief under pathogen, has been isolated from almost all the body site examined and the environment but Streptococci from fewer body sites, whereas the prevalence of Escherichia coli has been reported to be widespread [11], [12], [13]. Very limited research works on mastitis have been carried out in Bangladesh [14], [15], [16], [17]. For the purposes of the study, the author were very interested to know the prevalence of mastitis in dairy cow at Sylhet district.

## 2. Materials and Methods

### 2.1 Study areas

The study was conducted at 2 different areas of Sylhet district to study the prevalence of mastitis. The data was collected from UVH, Jaintapur Upazilla and District Dairy Farm, Sylhet.

### 2.2 Study population

Total cows of 49 were sampled from Veterinary Hospital of Jaintapur Upazilla and Dairy farm, Sylhet.

### 2.3 Period of study

The study on prevalence and use of antibiotics were conducted from 1<sup>st</sup> January 2016 to 31<sup>st</sup> December 2016 in the study area when stay at Veterinary Hospital of Jaintapur Upazilla and Dairy farm, Sylhet for internship base learning.

### 2.4 Source of data collection

Upazilla Veterinary Hospital and Dairy farm of Sylhet were the sources of population for conducting the study.

### 2.5 Methods of data collection

This data was collected from the record book of the Veterinary Hospital of Jaintapur Upazilla and Dairy farm, Sylhet. The following steps were taken for conducting the study. A structured questionnaire was developed and asked and the pre-tested questionnaire was filled. Each questionnaire asking time was about 10-15 minutes. The questionnaire contained information regarding age, breed, health, stage of lactation and management status. Open ended questions were collected and recorded.

### ➤ Moderate Form

Generalized swelling of infected quarter and painful udder was found. Abnormality in milk also was manifested without systemic reaction.

### ➤ Severe Form

The secretion was abnormal with hot and swollen quarter or udder. Fever, rapid pulse, loss of appetite, dehydration and depression was manifested.

### 2.8 Data Analysis

All collected data were entered into Microsoft Excel spreadsheet. The prevalence of clinical mastitis was the dependent variable while source of patient, age, general physical condition were independent variables considered at cow level.

### 3. Results

#### 3.1 Overall prevalence of mastitis

The data was collected from January 2016 to December 2016 at Jaintapur Upazilla Veterinary Hospital record book. In the month of January 2016, total 53 animals were examined in the Jaintapur Veterinary Hospital and from that 2 were affected as mastitis animal.

**Table 1:** Prevalence of mastitis of dairy cows from January 2016 to December 2016 at Jaintapur

Month	Animal examined	Affected animal ( Mastitis )	Prevalence ( % )
January	53	2	3.77
February	46	3	6.52
March	51	2	3.92
April	55	1	1.82
May	40	2	5
June	32	1	3.13
July	35	5	14.28
August	59	4	6.78
September	36	4	11.11
October	36	5	13.89
November	34	2	5.88
December	26	1	3.84
	Total-503	Total -36	7.15

So, the prevalence of mastitis in January was 3.77%. As the same way in February , March , April , May, June , July, August, September , October, November, December, the total number of animals examined were 46,51,55,40,32,35,59,36,36,34,26 respectively and the affected animals were 2,3,2,1,2,1,5,4,4,5,2,1. So the

prevalence of mastitis in those month were 6.52%, 3.92%, 1.82%, 5%, 3.13%, 14.28%, 6.78%, 11.11%, 13.89%, 5.88%, 3.84% respectively. From the above discussion it has been seen that the highest prevalence was in July and low in April. The overall prevalence of mastitis at Jaintapur was 7.15%.

**Table 2:** Prevalence of mastitis of dairy cows from January 2016 to December 2016 at dairy Farm, Sylhet

Month	Animal examined	Affected animal ( Mastitis )	Prevalence ( % )
January	30	1	3.33
February	14	0	0
March	11	1	9.09
April	19	1	5.26
May	18	1	5.55
June	16	0	0
July	8	2	25
August	16	1	6.25
September	11	2	18.18
October	12	2	16.67
November	12	1	8.33
December	21	2	9.52
	Total-188	Total-13	6.91

The data was collected from January 2016 to December 2016 at Dairy Farm record book. In the month of January 2016, total 30 animals were examined in the Jaintapur Veterinary Hospital and from that 1 was affected as mastitis animal. So, the prevalence of mastitis in January was 3.33%. As the same way in February , March , April , May, June , July, August, September , October, November, December, the total number of animals

examined were 14,11,19, 18,16,8,16,11,12,12,21 respectively and the affected animals were 0,1,1,1,0,2,1,2,2,1,2. So the prevalence of mastitis in those month were 0, 9.09%, 5.26%, 5.55%, 0, 25%, 6.25%, 18.18%, 16.67%, 8.33%, 9.52% respectively. From the above discussion it has been seen that the highest prevalence was in July and low in January. The overall prevalence of mastitis at Dairy Farm was 6.91%.

### 3.2 Prevalence according to age

**Table 3:**Prevalence of mastitis according to age at Jaintapur (2016)

Age (year)	Animal examined	Affected animal	Prevalence ( % )
3-6	167	8	4.79
7-8	141	10	7.09
9-19	195	18	9.23
	Total-503	Total-36	

Age is an important factor of mastitis .From table 3, it has been seen that the age between 9-19 years of animals were more susceptible to mastitis. So the prevalence of mastitis between 3-6 years was 4.79%. The prevalence of mastitis

between 7-8 years was 7.09%. The prevalence was 9.23%. From this data, the highest prevalence of mastitis is related to 9-19 years of age and the lowest prevalence was in 3-6 years of age.


**Table 4:**Prevalence of mastitis according to age at Dairy Farm, Sylhet (2016)

Age (year)	Animal examined	Affected animal	Prevalence ( % )
3-6	62	2	3.23
7-8	55	4	7.27
9-11	71	7	9.85
Total	188	13	

From table 4, it has been seen that the age between 9-19 years of animals were more susceptible to mastitis. From this data, the highest prevalence of mastitis was related to 9-19 years of age and the lowest prevalence was in 3-6 years of age.

The environment factor is one of the most important factors for the cause of mastitis in dairy cows. This data was collected from Jaintapur Upazilla Veterinary Hospital. From figure 1 , the prevalence of mastitis in summer season(March, April, May, June) was 3.47% and in rainy season ( July, August, September, October) was 11.52% and in winter season ( November, December, January, February) was 5%.

**3.3 Prevalence according to season**



**Figure 1.** Season wise Prevalence% of mastitis at Jaintapur Upazilla (2016) and Dairy Farm , Sylhet.

From the above discussion it has been seen that the highest prevalence of mastitis was in rainy season and low in summer season. From figure 1,the prevalence of mastitis at Dairy farm in summer season(March, April,

May, June) was 4.97% and in rainy season ( July, August, September, October) was 16.53% and in winter season (November, December, January, February) is 5.29% .

### 3.4 Breed related prevalence




Figure 2. Prevalence% of mastitis according to breed at Jaintapur Upazilla (2016)

The highest prevalence of mastitis was 7.19% found in local breed and lowest 6.79% in cross breed (Figure 2).

### 3.1.5 Lactation number-based prevalence of mastitis




Figure 3: Prevalence% of mastitis according to lactation number

The figure 3 showed that there was 2.89%, 5.18%, 8.33%, 8.75% and 15.38% prevalence at 1st, 2nd, 3rd, 4th and 5th lactation respectively. The rate of increasing prevalence was highly significant. The prevalence was much higher in 5th lactation than others.

## 4. Discussions

In Dairy Farm, Sylhet, the overall prevalence of mastitis was 8.24% and the prevalence of mastitis in Jaintapur was 2.20%. Findings of the present study was lower than the earlier findings of Kaderet al., [18] who reported 46.6%. In Jaintapur Veterinary Hospital, The prevalence of mastitis was recorded as 4.79%, 7.09%, and 9.23% respectively at the age group of 3-6 years, 7-8 years and 9-11 years. In Dairy Farm, Sylhet prevalence of mastitis was recorded as 3.23%, 7.27% and 9.85 % respectively at the age group of 3-6 years, 7-8 years and 9-11 years. The prevalence of clinical mastitis was also increased in older cows. Similar observation was also reported by Rahman et al., [19] and Pluvineget al., [20]. Rahman et al., [19] reported 57.5%

prevalence of mastitis in the age group of higher than 9 years old and 40.1% in the age group of 7 and 8 years. Similarly Biffa et al., [21] suggested older cows are at more risk (44.6%) for the incidence of mastitis than younger cows (23.6%). The present study revealed that high incidence of mastitis was recorded during monsoon season which is in agreement with Shinde et al., [22], Jadhav et al., [23] and Ameh et al. [24]. All the cows of Dairy Farm, Sylhet were cross breed and the prevalence of mastitis was 6.91% whereas local and cross breed of Jaintapur Upazilla, the prevalence of mastitis was 7.19% and 6.79% respectively. The results of this study was lower than that of Rabbani [25] who reported 20.31% in Holstein Friesian cross cows and 13.24% in Red Chittagong cows and of Prodhan et al., [17] who reported 16.52% and 15.77% incidence of mastitis. The exotic breeds like Jersey are more susceptible to bovine mastitis than indigenous breeds Dutta et al., [26] concluded the risk ratio of developing mastitis in Jersey was approximately two times higher than indigenous breeds. Similarly Biffa



et al., [21] reported HF cows are affected at higher rate (56.5%) compared with local zebu (30.9%) and Jersey cows (28.9%). The number of lactations had a significant effect on the prevalence of mastitis irrespective of seasons. The prevalence of mastitis was the lowest in first lactation and the highest in 5<sup>th</sup> lactation. Regarding parity, several studies were in agreement with the present findings of increased mastitis in advancing age and parity [27].

## 5. Conclusion

The overall prevalence of clinical mastitis in cows was relatively lower in the study area. The prevalence of mastitis was higher in advancing age and in 5<sup>th</sup> lactation period. Mastitis is one of the major important diseases of milch cows throughout the world including Bangladesh. Proper hygienic management of the farm house and utensils and treatment regimens should be followed to reduce the economic loss and also the prevalence of disease different epidemiological factors that interplay in mastitis occurrence.

## 6. Recommendations

- Strict maintain hygienic measurement during milking time.
- Post-milking teat dipping by antiseptics.
- Dry cow treatment of all quarters of all cows.
- Use of properly functioning milking machines.
- Culling of chronic mastitis cows from the farm.
- Use of segregation if practical.
- Treatment of new clinical cases promptly.

## 7. Acknowledgement

This study was the Internship report writing of the 1<sup>st</sup> author. The author was very much grateful to the authority of dairy farm, Sylhet and veterinary Surgeon, Jaintapur Upazilla for providing all necessary information to complete this study. All authors were share their scientific knowledge in optimum level and approved the final manuscript. It was an internship report writing of 1<sup>st</sup> author.

## 8. References

- [1]. DLS. Annual report on livestock. Division of Livestock Statistics, Ministry of Fisheries and Livestock, Farmgate, Dhaka, Bangladesh, 2008.
- [2]. T. Hemme, "IFCN Dairy Report". International Farm Comparison Network. IFCN Dairy Research Center. Kiel Germany, 2008.
- [3]. FAO. Food and Agriculture Organization of the United Nations. Ministry of Fisheries and Livestock. Bangladesh, pp. 65, 2008:
- [4]. B.k. Devi, P.C. Shukla and R.K. Bagherwal, "Incidence of sub-clinical mastitis in cows", *Indian Journal Dairy Science*, vol. 50, pp. 477-478, 1997.
- [5]. W.N. Philpot, "Economics of mastitis control", *Veterinary Clinics of North America Large Animal Practice*, vol. 6, pp. 233 – 235, 1984.
- [6]. O.M. Radostits, D.C. Blood and C.C. Gay, "Veterinary Medicine : A textbook of the disease of cattle, sheep, pigs, goat , horses", 8<sup>th</sup>Edn. BacilieriTindall, pp. 563- 616, 1994.
- [7]. M. Itagaki, S. Abe, J. Sakai and K. Suzuki, "Relationship between morphologic abnormalities of teat orifice and subclinical mastitis in dairy cattle", *Journal of Japan Veterinary Medical Association*, vol. 52, pp.561 – 564,1999.
- [8]. B.B. Malhotra and M.P. Kapur, "Epizootiological studies on bacterial bovine mastitis", *Indian Veterinary Journal*, vol. 59, pp 921 – 926, 1982.
- [9]. Prabhakar S K, Singh K B, Nauriyal D C and Sidhu S S. Epizootiological studies of mastitis causing organisms in cross-bred cows. *Indian Veterinary Journal*, 1990; 67: 734 – 738.
- [10]. B.Pal, B.B. Verma and R.S.A. Prasad, "A note on incidence of subclinical mastitis in cows and buffaloes in an organized farm at Ranchi", *Indian Journal of Veterinary Medicine*, vol.11, pp. 32 – 33, 1991.
- [11]. H.D. Joshi, B.R. Joshi and H.K. Shrestha, "Epidemiological investigation on clinical mastitis in cattle and buffaloes in the Western hills of Nepal", *Veterinary Review*, vol. 13, pp.12 – 15, 1998.
- [12]. M.M. Sen, M.H. Kabir and A. Rahman, "Application of indirect tests to detect subclinical mastitis in milch cows", *Bangladesh Veterinary Journal*, vol. 30, pp. 137 – 139, 1996.
- [13]. M.S. Rahman, M. Nooruddin and M.M. Rahman, "Prevalence and distribution of mastitis in crossbred and exotic dairy cows", *Bangladesh Veterinarian*, vol. 14, pp. 1-4, 1997.
- [14]. M.A.M. Prodhana, A.H.M. Kamal and A.T.M. Mahbub-E-Elahi, "Prevalence of subclinical mastitis in cows of Baghabari Milk-shed Area", *Bangladesh Veterinary Journal*, vol. 130, pp. 59-61, 1993.
- [15]. M.A. Kader, M.A. Samad and S. Saha, "Influence of host level factors on prevalence and economics of sub-clinical mastitis in dairy cows in Bangladesh", *Indian Journal of Dairy Science*, vol.56, pp 235-240, 2003.
- [16]. P. Pluvinaige, T. Ducruet, J. Josse and F. Monicat, "Factors of risk of milk cows mastitis, Skara, Sweeden", *Environment and Animal Hygiene*, vol.1, pp. 51-55, 1990.
- [17]. D. Biffa, E. Debela and F. Beyene, "Prevalence and risk factors of mastitis in lactating dairy cows in Southern Ethiopia", *International Journal of*

Applied Research in Veterinary Medicine, vol. 3(3), pp.189–198, 2005.

- [18]. S.S. Shinde, G.B. Kulkarni, G.K. Gangane and N.M.D. Degloorkar, "Incidence of mastitis in Proceedings of the Round Table Conference on prebhani district", Maharashtra, Ludhiana, India, 2001.
- [19]. K.L. Jadhav, V.N. Tripathi and M.M. Kale, "Incidence and economics and mammary disorders in Holstein X Sahiwalcross bred cows", Indian Journal of dairy Science, vol. 48, pp.382–385, 1995.
- [20]. A.F.M.G. Rabbani and M.A. Samad, "Host determinants based comparative prevalence of subclinical mastitis in lactating Holstein-Frisian cross cows and Red Chittagong cows in Bangladesh", Bangladesh Journal of Veterinary Medicine, vol. 8(1), pp. 17-21, 2008.
- [21]. J. Dutta, B.S. Rathore and S.G. Mullick, "Lactational incidence rate of mastitis in exotic and crossbreed cows, An epidemiology study", Indian Journal of Veterinary Pathology, vol.12, pp. 28–32, 1990.
- [22]. M.A. Quaderi, "Prevalence of sub-clinical mastitis in dairy farms". MS Thesis, Department of Surgery and Obstetrics, Faculty of Veterinary Science, Bangladesh Agricultural University, Mymensingh, Bangladesh, 2005..



Md Mowdudul Hasan Talha completed his DVM degree at 2017 from Faculty of Veterinary, Animal and Biomedical Sciences, Sylhet Agricultural University with scholastic and outstanding result and stood 2nd position among all in his passing year.

Now currently he is doing his master's in Department of Pharmacology and Toxicology in Sylhet Agricultural University. He is highly interested in wildlife conservation and eco health approach of wild animals.



Mamun Ur Rahman, completed DVM degree from Sylhet Agricultural University. Now he is doing his post graduation course in the subject of genetics and animal breeding. Besides this he is working on the productive performance of duck in sylhet

region.



Mustaq Ahmad, completed DVM degree from Sylhet Agricultural University. Now he is doing his post graduation course at genetics and animal breeding department. Besides this he is working on the productive performance of duck in sylhet region.



Akash Khasnobish, completed DVM degree from Sylhet Agricultural University. Now currently doing his master's in Department of Pharmacology and Toxicology in Sylhet Agricultural University.



Md Abdus Sabur, completed DVM degree from Sylhet Agricultural University. Now he is doing his master's in Department of Medicine. He has high interest for finding the efficacy of different types of antibiotic in case of Mastitis.



Md Altafur Rahman currently enrolled his MS in genetics and animal breeding under the department of genetics and animal breeding. His key interest is to find interrelations between drugs and genetic changes among animals with their morphometric traits. He is currently doing research about

morphometric characters of red jungle fowl.

## Authors Profile



Jannatul Mawa Momu, an enthusiastic fellow of veterinary sector of Bangladesh has completed her DVM degree in 2017 from Faculty of Veterinary, Animal and Biomedical Sciences, Sylhet Agricultural University, stood 1st position among her session with brilliant result.

Currently she is doing her post graduation at Department of Animal Nutrition, SAU. She is very interested to work with rumenflora and other digestive beneficial microflora, micronutrient and invtro mineralization.



Mahfuz Rahman Adnan, a newly introduced fellow of veterinary sector of Bangladesh has completed his DVM degree in 2017 from Faculty of Veterinary, Animal and Biomedical Sciences of Sylhet Agricultural University. Now

currently doing his master's in pathology in Sylhet Agricultural University. He wants to work on molecular research about Avian influenza and other emmerging diseases.



Dr. MD. Matiur Rahman is working as an Assistant Professor, Department of Medicine, Faculty of Veterinary, Animal and Biomedical Sciences, Sylhet Agricultural University, Bangladesh. He has completed his graduation in Doctor of

Veterinary Medicine at 2013, securing CGPA 3.887 out of 4.00 and stood first position with distinction and obtained his Master of Science in Medicine degree from the same institution, securing CGPA 4.0 out of 4.0 and stood first position with distinction. During his undergraduate program the areas that evoked interest in was “biomedical research” and inspired to pursue MS in Medicine. He got Prime Minister Gold Medal Awaed-2013 for his brilliant result in his under garduate studies. He had strong desire to develop his skills especially in innovative biomedical technologies. He has published 17 research articles in established and renowned national and international scientific journals. His research interest on bovine cancer, proteomics and infectious and non infectious diseasees of birds espicially pigeon.